Math 1B Section 107 Quiz #7

Thursday, 11 October 2007

GSI: Theo Johnson-Freyd
http://math.berkeley.edu/∼theojf

Name: ____

1. **True or False** (1 pt each) For each of the following statements, decide if it is true or false. You do not need to show work: I will grade only your answers.

   (a) Let's say $0 \leq a_n \leq b_n$, and $\sum_{n=1}^{\infty} a_n = A$ and $\sum_{n=1}^{\infty} b_n = B$. If $A = B$, then $a_n = b_n$ for every n.
   TRUE. The first inequality guarantees that $\sum(b_n - a_n) = B - A = 0$ is a sum of nonnegative terms, but the only way to add up nonnegative numbers and get 0 is if each of the numbers is itself 0.

   (b) If $0 \leq a_n \leq f(n)$, where $f(x)$ is a continuous decreasing function on $x \in [1, \infty)$, such that $\int_1^{\infty} f(x)\, dx$ converges, then $\sum_{n=1}^{\infty} a_n$ converges.
   TRUE. This is a combination of the comparison and integral tests.

   (c) If $0 \leq a_n \leq \pi/2$ and $\sum_{n=1}^{\infty} \sin(a_n)$ diverges, then $\sum_{n=1}^{\infty} a_n$ diverges.
   TRUE. This follows from the comparison test, since $\sin(x) < x$ if $x > 0$, and so $a_n > \sin(a_n)$.

For the next two questions, use either the **Limit Comparison Test** or the **Integral Test** to determine if the series converges or diverges. Be sure to check that the series satisfies the conditions necessary for the test.

2. (3 pts) $\displaystyle\sum_{n=1}^{\infty} \frac{\arctan(n)}{n^2 - \ln(n)}$

   We recall that $\lim_{n\to\infty} \arctan(n) = \pi/2$*. Then comparing with* $1/n^2$ *(which converges) gives*

   $$\lim_{n\to\infty} \frac{\arctan(n)/(n^2 - \ln n)}{1/n^2} = \lim_{n\to\infty} \arctan(n) \frac{1}{1 - (\ln(n)/n^2)} = \frac{\pi}{2} \cdot 1$$

   which is more than 0 *and less than* ∞*. So since* $\sum 1/n^2$ *converges, so must our series* $\sum \arctan(n)/(n^2 - \ln n)$*.*

3. (4 pts) $\displaystyle\sum_{n=3}^{\infty} \frac{1}{n \ln n \ln(\ln n)}$

   Recognizing this as something integrable, we check the conditions for the integral test:

   - $1/(x \ln x \ln(\ln x))$ *is always positive.*
   - *It is the product of three decreasing functions* ($1/x$, $1/\ln x$, *and* $1/\ln\ln x$)*, so must itself be decreasing.*

   Thus, using the integral test, we check

   $$\int_3^{\infty} \frac{dx}{x \ln x \ln(\ln x)} = \int_{\ln\ln 3}^{\infty} \frac{du}{u} = \infty$$

   where we substitute $u = \ln(\ln x)$*. Thus the original series also diverges.*