

Math 1B Worksheet 4: Integration by Parts

Thursday, 6 September 2007

Theo Johnson-Freyd
theo.jf@math.berkeley.edu

Please introduce yourselves to each other, and put your names at the top of a piece of blackboard. Take turns being the scribe: each of you should have a chance to write on the chalkboard for at least one of the exercises.

These exercises are hard — harder than on the homework, quizzes, or exams. That means that you should spend some time thinking and talking about them; they're designed to be solved in groups (the best way to learn mathematics). The problems are roughly in order of increasing difficulty. I don't expect any group to solve all of them.

Don't forget to draw pictures.

1. Integrate:

$$\int \frac{dx}{3 \cos x + 4 \sin x}$$

2. Let's find the partial fraction decomposition of, say,

$$\frac{x+1}{x^3+x^2-2x}$$

- (a) Factor the denominator to find a , b , and c :

$$\frac{x+1}{x^3+x^2-2x} = \frac{x+1}{(x-a)(x-b)(x-c)}$$

- (b) We want to write

$$\frac{x+1}{x^3+x^2-2x} = \frac{A}{x-a} + \frac{B}{x-b} + \frac{C}{x-c}$$

We could put everything over a common denominator (i.e. multiply by (x^3+x^2-2x)) and compare like terms. Instead, multiply both sides by just $(x-a)$, and then plug in $x=a$ into both sides of the equation. What does this tell you about the values of A , B , and C ?

- (c) Use this method to find A , B , and C .
 (d) How would you do this for other examples? What about

$$\frac{x+1}{x^3-2x^2+x} ? \quad \frac{x+1}{x^3-x^2+x} ?$$

3. Solve

$$\int \frac{x dx}{x^2-1}$$

in three different ways:

- (a) With a u -substitution.
 (b) With a trig substitution.
 (c) By decomposing the integrand into partial fractions.

For an extra challenge, try doing it by parts: $u = x$, $dv = dx/(x^2 - 1)$, and integrate dv to find v with whatever method you want.

4. When a marble (with mass m , say) falls through a viscous liquid like honey, a constant downward force (gravity minus buoyancy = $\tilde{g} = mg - b$) acts on it, and friction impedes its motion with a force proportional to the square of the marble's velocity (say αv^2). Then the marble's velocity is given by the equation

$$\int dt = \int \frac{m dv}{\tilde{g} - \alpha v^2}$$

To simplify the problem, let's let $m = \tilde{g} = \alpha = 1$ (or, if you want to, do the problem with all the unknown constants). Assume that the marble starts at rest at time $t = 0$. Solve this integral to find v as a function of t ; don't forget to use the fact that $v(t = 0) = 0$ to figure out the constant of integration.

For an extra challenge, find the distance the marble has fallen as a function of t .

5. **This problem is only for those who know complex numbers.**

Normally we integrate

$$\int \frac{dx}{x^2+1}$$

by a trig substitution. Try factoring the denominator with complex numbers and using integration by parts.