

Math 53 Quiz 12

9 May 2008

GSI: Theo Johnson-Freyd
<http://math.berkeley.edu/~theo/f/>

Name: _____

Time (circle one):

12:10 - 1:00

3:10 - 4:00

Please use extra paper as necessary. For each part, partial credit will be assigned based on correct work (you do need to show some work, enough so that I know how you solved the problem). Please simplify and box your answers.

When a solid body is partially submerged in liquid, it feels a water-pressure force on each point of its surface. If the water pressure at a point (x, y, z) is $P(x, y, z)$, then the force on a small area dS with outward normal $\hat{\mathbf{n}}$ of the surface is:

$$\frac{d\vec{\mathbf{F}}}{dS} = -P \hat{\mathbf{n}}$$

(The minus sign is because the pressure points in.) The aggregate of all the little bits of pressure can act as a buoyancy force; the total force is:

$$\vec{\mathbf{F}}_{\text{total}} = - \iint_S P \hat{\mathbf{n}} dS$$

where S is the surface of the object under the water. The component of $\vec{\mathbf{F}}$ in the $\hat{\mathbf{i}}$ -direction (or $\hat{\mathbf{j}}$ or $\hat{\mathbf{k}}$ or anything else) is:

$$F_{\hat{\mathbf{i}}} = \hat{\mathbf{i}} \cdot \vec{\mathbf{F}} = \iint_S -P \hat{\mathbf{i}} \cdot \hat{\mathbf{n}} dS$$

In general, the pressure is

$$P(x, y, z) = \begin{cases} -\rho g z, & z \leq 0 \\ 0, & z \geq 0 \end{cases}$$

where ρ and g are constants (physically, ρ is the density of water, and g is the strength of gravity); the minus sign is because we measure z going up, but pressure is higher as you go down. We set $P(x, y, z) = 0$ for $z \geq 0$, thinking of the water level as $z = 0$, and pretending that there is no air pressure.

- a. (3 pts) Is $-P\hat{\mathbf{i}} = \rho g z \hat{\mathbf{i}}$ the curl of a vector field? If not, explain why not. If so, find a vector field $\vec{\mathbf{v}}$ so that $\vec{\nabla} \times \vec{\mathbf{v}} = -P\hat{\mathbf{i}}$.

- b. (2 pts) Consider the situation when the object is only partially submerged, as in Figure 1. Let S be the surface below the level of the water, and V the volume of the object below the water. Calculate, in terms of V , ρ , and g , the integral:

$$F_{\hat{\mathbf{i}}} = \iint_S -P\hat{\mathbf{i}} \cdot \hat{\mathbf{n}} dS = \iint_S \rho g z \hat{\mathbf{i}} \cdot \hat{\mathbf{n}} dS$$

- c. (3 pts) Is $-P\hat{\mathbf{k}} = \rho g z \hat{\mathbf{k}}$ the curl of a vector field? If not, explain why not. If so, find a vector field $\vec{\mathbf{v}}$ so that $\vec{\nabla} \times \vec{\mathbf{v}} = -P\hat{\mathbf{k}}$.

- d. (2 pts) Consider the situation when the object is fully submerged, as in Figure 2. Let S be the surface below the level of the water, and V the volume of the object below the water. Calculate, in terms of V , ρ , and g , the integral:

$$F_{\hat{\mathbf{k}}} = \iint_S -P\hat{\mathbf{k}} \cdot \hat{\mathbf{n}} dS = \iint_S \rho g z \hat{\mathbf{k}} \cdot \hat{\mathbf{n}} dS$$